

Fall 2022

INTRODUCTION TO COMPUTER VISION

Atlas Wang

Assistant Professor, The University of Texas at Austin

Visual Informatics Group@UT Austin


<https://vita-group.github.io/>



Camera Model

Pinhole and Lens

Pinhole camera a.k.a. camera obscura




Pinhole camera a.k.a. camera obscura

First mention ...



Chinese philosopher Mozi
(470 to 390 BC)

First camera ...



Greek philosopher Aristotle
(384 to 322 BC)

Pinhole camera terms

real-world
object



barrier (diaphragm)



pinhole
(aperture)



digital sensor
(CCD or CMOS)

Pinhole camera terms

real-world
object



barrier (diaphragm)



pinhole
(aperture)



camera center
(center of projection)


image plane



digital sensor
(CCD or CMOS)

Focal length

real-world
object




focal length f

Focal length

What happens as we change the focal length?


real-world
object



Focal length

What happens as we change the focal length?

real-world
object




Focal length

What happens as we change the focal length?

object projection is half the size

real-world
object



Pinhole size

real-world
object



pinhole
diameter



Ideal pinhole has infinitesimally small size

- In practice that is impossible.

Pinhole size

What happens as we change the pinhole diameter?

real-world
object




pinhole
diameter



Pinhole size

What happens as we change the pinhole diameter?


real-world
object



Pinhole size

What happens as we change the pinhole diameter?

real-world
object




Pinhole size

What happens as we change the pinhole diameter?


object projection becomes blurrier

real-world
object



Extreme Case: Bare-sensor imaging

real-world
object



digital sensor
(CCD or CMOS)

All scene points contribute to all sensor pixels

What does the
image on the
sensor look like?

Extreme Case: Bare-sensor imaging




All scene points contribute to all sensor pixels

What about light efficiency?

real-world
object




pinhole
diameter



focal length f

- What is the effect of doubling the pinhole diameter?
- What is the effect of doubling the focal length?


The lens camera




Lenses map “bundles” of rays from points on the scene to the sensor.

How does this mapping work exactly?

The pinhole camera




The lens camera



Central rays propagate in the same way for both models!


Important difference: focal length

In a pinhole camera, focal length is distance between aperture and sensor





Important difference: focal length

In a lens camera, focal length is distance where parallel rays intersect



Describing both lens and pinhole cameras



We can derive properties and descriptions that hold for both camera models if:

- We use only central rays.
- We assume the lens camera is in focus.
- We assume that the focus distance of the lens camera is equal to the focal length of the pinhole camera.

Remember: *focal length* f refers to different things for lens and pinhole cameras.

- In this lecture, we use it to refer to the aperture-sensor distance, as in the pinhole camera case.

Camera Matrix

The camera as a coordinate transformation


The camera as a coordinate transformation

A camera is a mapping from:

the 3D world

to:

a 2D image



The camera as a coordinate transformation

A camera is a mapping from:

the 3D world

to:

a 2D image

homogeneous coordinates

$$\mathbf{x} = \mathbf{P}\mathbf{X}$$

2D image
point

camera
matrix

3D world
point

What are the dimensions of each variable?

The camera as a coordinate transformation

$$\mathbf{x} = \mathbf{P}\mathbf{X}$$


$$\begin{bmatrix} X \\ Y \\ Z \end{bmatrix} = \begin{bmatrix} p_1 & p_2 & p_3 & p_4 \\ p_5 & p_6 & p_7 & p_8 \\ p_9 & p_{10} & p_{11} & p_{12} \end{bmatrix} \begin{bmatrix} X \\ Y \\ Z \\ 1 \end{bmatrix}$$

homogeneous
image coordinates
3 x 1

camera
matrix
3 x 4

homogeneous
world coordinates
4 x 1


The pinhole camera




The (rearranged) pinhole camera

image plane

real-world
object




The (rearranged) pinhole camera




What is the equation for image coordinate \mathbf{x} in terms of \mathbf{X} ?

The 2D view of the (rearranged) pinhole camera




What is the equation for image coordinate x in terms of X ?

The 2D view of the (rearranged) pinhole camera



The (rearranged) pinhole camera



What is the camera matrix \mathbf{P} for a pinhole camera?

$$\mathbf{x} = \mathbf{P}\mathbf{X}$$

Homogeneous Coordinates

Given a point \mathbf{p} in \mathbb{R}^2 , represented as $P = (p_1, p_2)$, i.e., the vector $\mathbf{p} = [p_1 \ p_2]^T$ its homogeneous representation (using homogeneous coordinates) is

$$\tilde{\mathbf{p}} = [\tilde{p}_1 \ \tilde{p}_2 \ \tilde{p}_3]^T; \quad \text{with } [0 \ 0 \ 0]^T \text{ not allowed}$$

The vector representation is obtained dividing the first n homogeneous components by the $(n + 1)$ -th, that is often called **scale**.

$$p_1 = \tilde{p}_1 / \tilde{p}_3; \quad p_2 = \tilde{p}_2 / \tilde{p}_3$$




Figure: Geometric interpretation of homogeneous coordinates.

Take-Away:

- All points, on the same projection ray, are mapped to the same homogeneous coordinate!
- It simplifies many equations in projective geometry! **Let's see next page...**

The pinhole camera matrix

Relationship from similar triangles:

$$[X \quad Y \quad Z]^T \rightarrow [X/Z \quad Y/Z]$$

Normal coordinates:

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} X/Z \\ Y/Z \end{bmatrix}$$

General camera model in *homogeneous coordinates*:

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} p_1 & p_2 & p_3 & p_4 \\ p_5 & p_6 & p_7 & p_8 \\ p_9 & p_{10} & p_{11} & p_{12} \end{bmatrix} \begin{bmatrix} X \\ Y \\ Z \\ 1 \end{bmatrix}$$

Homogenous coordinates:

$$\begin{bmatrix} x \\ y \\ z (= 1) \end{bmatrix} =$$

What does the pinhole camera projection look like?

$$\mathbf{P} = \begin{bmatrix} ? & ? & ? & ? \\ ? & ? & ? & ? \\ ? & ? & ? & ? \end{bmatrix}$$

$$\begin{bmatrix} X/Z \\ Y/Z \\ 1 \end{bmatrix} = \begin{bmatrix} X \\ Y \\ Z \end{bmatrix}$$

The pinhole camera matrix

Relationship from similar triangles:

$$[X \quad Y \quad Z]^T \rightarrow [X/Z \quad Y/Z]$$

Normal coordinates:

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} X/Z \\ Y/Z \end{bmatrix}$$

General camera model *in homogeneous coordinates*:

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} p_1 & p_2 & p_3 & p_4 \\ p_5 & p_6 & p_7 & p_8 \\ p_9 & p_{10} & p_{11} & p_{12} \end{bmatrix} \begin{bmatrix} X \\ Y \\ Z \\ 1 \end{bmatrix}$$

Homogenous coordinates:

$$\begin{bmatrix} x \\ y \\ z (= 1) \end{bmatrix} =$$

What does the pinhole camera projection look like?

The perspective projection matrix

$$\mathbf{P} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

$$\begin{bmatrix} X/Z \\ Y/Z \\ 1 \end{bmatrix} = \begin{bmatrix} X \\ Y \\ Z \end{bmatrix}$$

The pinhole camera matrix

Relationship from similar triangles:

$$[X \quad Y \quad Z]^T \rightarrow [X/Z \quad Y/Z]$$

General camera model *in homogeneous coordinates*:

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} p_1 & p_2 & p_3 & p_4 \\ p_5 & p_6 & p_7 & p_8 \\ p_9 & p_{10} & p_{11} & p_{12} \end{bmatrix} \begin{bmatrix} X \\ Y \\ Z \\ 1 \end{bmatrix}$$


What does the pinhole camera projection look like?

The perspective
projection matrix

$$\mathbf{P} = \begin{bmatrix} 1 & 0 & 0 & | & 0 \\ 0 & 1 & 0 & | & 0 \\ 0 & 0 & 1 & | & 0 \end{bmatrix} = [\mathbf{I} \quad | \quad \mathbf{0}]$$

alternative way to write
the same thing


More general case: arbitrary focal length



What is the camera matrix \mathbf{P} for a pinhole camera?


$$\mathbf{x} = \mathbf{P}\mathbf{X}$$

More general (2D) case: arbitrary focal length



What is the equation for image coordinate x in terms of X ?

More general (2D) case: arbitrary focal length



$$[X \ Y \ Z]^T \mapsto [fX/Z \ fY/Z]^T$$

The pinhole camera matrix for arbitrary focal length

Relationship from similar triangles:

$$[X \quad Y \quad Z]^T \mapsto [fX/Z \quad fY/Z]^T$$

General camera model *in homogeneous coordinates*:


$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} p_1 & p_2 & p_3 & p_4 \\ p_5 & p_6 & p_7 & p_8 \\ p_9 & p_{10} & p_{11} & p_{12} \end{bmatrix} \begin{bmatrix} X \\ Y \\ Z \\ 1 \end{bmatrix}$$

What does the pinhole camera projection look like?

$$\mathbf{P} = \begin{bmatrix} f & 0 & 0 & 0 \\ 0 & f & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$


Generalizing the camera matrix

In general, the camera and image have *different* coordinate systems.



Generalizing the camera matrix

In particular, the camera origin and image origin may be different:




$$\begin{bmatrix} X \\ Y \\ Z \end{bmatrix} = \begin{bmatrix} p_1 & p_2 & p_3 & p_4 \\ p_5 & p_6 & p_7 & p_8 \\ p_9 & p_{10} & p_{11} & p_{12} \end{bmatrix} \begin{bmatrix} X \\ Y \\ Z \\ 1 \end{bmatrix}$$

How does the camera matrix change?

$$\mathbf{P} = \begin{bmatrix} f & 0 & 0 & 0 \\ 0 & f & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

Generalizing the camera matrix

In particular, the camera origin and image origin may be different:



$$\begin{bmatrix} X \\ Y \\ Z \end{bmatrix} = \begin{bmatrix} p_1 & p_2 & p_3 & p_4 \\ p_5 & p_6 & p_7 & p_8 \\ p_9 & p_{10} & p_{11} & p_{12} \end{bmatrix} \begin{bmatrix} X \\ Y \\ Z \\ 1 \end{bmatrix}$$

How does the camera matrix change?

$$\mathbf{P} = \begin{bmatrix} f & 0 & p_x & 0 \\ 0 & f & p_y & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

shift vector
transforming
camera origin to
image origin

Camera matrix decomposition

We can decompose the camera matrix like this:

$$\mathbf{P} = \begin{bmatrix} f & 0 & p_x \\ 0 & f & p_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 & | & 0 \\ 0 & 1 & 0 & | & 0 \\ 0 & 0 & 1 & | & 0 \end{bmatrix}$$

What does each part of the matrix represent?

Camera matrix decomposition

We can decompose the camera matrix like this:

$$\mathbf{P} = \begin{bmatrix} f & 0 & p_x \\ 0 & f & p_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 & | & 0 \\ 0 & 1 & 0 & | & 0 \\ 0 & 0 & 1 & | & 0 \end{bmatrix}$$


↑
(homogeneous) transformation
from 2D to 2D, accounting for not
unit focal length and origin shift

↑
(homogeneous) perspective projection
from 3D to 2D, assuming **image plane at**
 $z = 1$ and **shared camera/image origin**

Also written as: $\mathbf{P} = \mathbf{K}[\mathbf{I}|\mathbf{0}]$ where $\mathbf{K} = \begin{bmatrix} f & 0 & p_x \\ 0 & f & p_y \\ 0 & 0 & 1 \end{bmatrix}$



Generalizing the camera matrix

In general, there are *three*, generally different, coordinate systems.




We need to know the transformations between them.

World-to-camera coordinate system transformation




tilde means
heterogeneous
coordinates

World-to-camera coordinate system transformation




World-to-camera coordinate system transformation



$$(\tilde{X}_w - \tilde{C})$$

translate

World-to-camera coordinate system transformation



$$R \cdot (\tilde{X}_w - \tilde{C})$$

rotate translate

Modeling the coordinate system transformation

In heterogeneous coordinates, we have:

$$\tilde{\mathbf{X}}_c = \mathbf{R} \cdot (\tilde{\mathbf{X}}_w - \tilde{\mathbf{C}})$$

How do we write this transformation in homogeneous coordinates?

Modeling the coordinate system transformation

In heterogeneous coordinates, we have:

$$\tilde{\mathbf{X}}_c = \mathbf{R} \cdot (\tilde{\mathbf{X}}_w - \tilde{\mathbf{C}})$$

In homogeneous coordinates, we have: (pay attention to R and C dimension!)

$$\begin{bmatrix} X_c \\ Y_c \\ Z_c \\ 1 \end{bmatrix} = \begin{bmatrix} \mathbf{R} & -\mathbf{RC} \\ \mathbf{0} & 1 \end{bmatrix} \begin{bmatrix} X_w \\ Y_w \\ Z_w \\ 1 \end{bmatrix} \quad \text{or} \quad \mathbf{X}_c = \begin{bmatrix} \mathbf{R} & -\mathbf{RC} \\ \mathbf{0} & 1 \end{bmatrix} \mathbf{X}_w$$

Incorporating the transform in the camera matrix

The previous camera matrix is for homogeneous 3D coordinates in camera coordinate system:

$$\mathbf{x} = \mathbf{P}\mathbf{X}_c = \mathbf{K}[\mathbf{I}|\mathbf{0}]\mathbf{X}_c$$

We also just derived:

$$\mathbf{X}_c = \begin{bmatrix} \mathbf{R} & -\mathbf{R}\tilde{\mathbf{C}} \\ \mathbf{0} & 1 \end{bmatrix} \mathbf{X}_w$$

Putting it all together

We can write everything into a single projection:

$$\mathbf{x} = \mathbf{P}\mathbf{X}_w$$

The camera matrix now looks like:

$$\mathbf{P} = \begin{bmatrix} f & 0 & p_x \\ 0 & f & p_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \mathbf{I} & | & \mathbf{0} \end{bmatrix} \begin{bmatrix} \mathbf{R} & -\mathbf{R}\tilde{\mathbf{C}} \\ \mathbf{0} & 1 \end{bmatrix}$$

intrinsic parameters (3 x 3):
correspond to camera
internals (**image-to-image**
transformation)

perspective projection (3 x 4):
maps 3D to 2D points
(**camera-to-image**
transformation)

extrinsic parameters (4 x 4):
correspond to camera
externals (**world-to-camera**
transformation)

Putting it all together


We can write everything into a single projection:

$$\mathbf{x} = \mathbf{P}\mathbf{X}_w$$


The camera matrix now looks like:

$$\mathbf{P} = \begin{bmatrix} f & 0 & p_x \\ 0 & f & p_y \\ 0 & 0 & 1 \end{bmatrix} \left[\mathbf{R} \quad \vdots \quad -\mathbf{RC} \right]$$

intrinsic parameters (3 x 3):
correspond to camera internals
(sensor not at $f = 1$ and origin shift)



extrinsic parameters (3 x 4):
correspond to camera externals
(world-to-image transformation)



General pinhole camera matrix

$$\mathbf{P} = \mathbf{K}[\mathbf{R}|\mathbf{t}] \quad \text{Or:} \quad \mathbf{P} = \mathbf{KR}[\mathbf{I} | -\mathbf{C}]$$

$$\mathbf{P} = \begin{bmatrix} f & 0 & p_x \\ 0 & f & p_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} r_1 & r_2 & r_3 & \cdots & t_1 \\ r_4 & r_5 & r_6 & \cdots & t_2 \\ r_7 & r_8 & r_9 & \cdots & t_3 \end{bmatrix}$$

intrinsic
parameters

extrinsic
parameters

$$\mathbf{R} = \begin{bmatrix} r_1 & r_2 & r_3 \\ r_4 & r_5 & r_6 \\ r_7 & r_8 & r_9 \end{bmatrix} \quad \mathbf{t} = \begin{bmatrix} t_1 \\ t_2 \\ t_3 \end{bmatrix}$$

3D rotation

3D translation

Recap

What is the size and meaning of each term in the camera matrix?

$$\mathbf{P} = \mathbf{KR}[\mathbf{I} \mid -\mathbf{C}]$$

The diagram shows the equation $\mathbf{P} = \mathbf{KR}[\mathbf{I} \mid -\mathbf{C}]$ with four arrows pointing upwards from question marks to the terms \mathbf{P} , \mathbf{KR} , $[\mathbf{I} \mid -\mathbf{C}]$, and \mathbf{C} . The arrows originate from question marks located below the terms: one under \mathbf{P} , one under \mathbf{KR} , one under the vertical bar in $[\mathbf{I} \mid -\mathbf{C}]$, and one under \mathbf{C} .

Recap

What is the size and meaning of each term in the camera matrix?

$$\mathbf{P} = \mathbf{KR}[\mathbf{I} | \mathbf{C}]$$

The diagram shows the equation $\mathbf{P} = \mathbf{KR}[\mathbf{I} | \mathbf{C}]$ with four arrows pointing from labels below to the terms \mathbf{P} , \mathbf{K} , \mathbf{I} , and \mathbf{C} . The label for \mathbf{P} is "3x3" and "intrinsic". The labels for \mathbf{K} , \mathbf{I} , and \mathbf{C} are each a question mark.

3x3
intrinsic

?

?

?

Recap

What is the size and meaning of each term in the camera matrix?

$$\mathbf{P} = \mathbf{KR}[\mathbf{I} | \mathbf{C}]$$

The diagram shows the equation $\mathbf{P} = \mathbf{KR}[\mathbf{I} | \mathbf{C}]$ with four arrows pointing from labels below to the terms \mathbf{K} , \mathbf{R} , $[\mathbf{I} |$, and $\mathbf{C}]$. The label '3x3 intrinsics' is positioned below \mathbf{K} , '3x3 3D rotation' is below \mathbf{R} , and two question marks are below the $[\mathbf{I} |$ and $\mathbf{C}]$ terms respectively.

3x3
intrinsics

3x3
3D rotation

?

?

Recap

What is the size and meaning of each term in the camera matrix?

$$\mathbf{P} = \mathbf{KR}[\mathbf{I} | \mathbf{C}]$$

The diagram shows the equation $\mathbf{P} = \mathbf{KR}[\mathbf{I} | \mathbf{C}]$ with four arrows pointing from labels below to the terms \mathbf{K} , \mathbf{R} , $[\mathbf{I} |$, and $\mathbf{C}]$. The labels are: 3×3 intrinsics, 3×3 3D rotation, 3×3 identity, and a question mark.

3×3 intrinsics 3×3 3D rotation 3×3 identity ?

Recap

What is the size and meaning of each term in the camera matrix?

$$\mathbf{P} = \mathbf{K}\mathbf{R}[\mathbf{I} \mid \mathbf{C}]$$

The diagram shows the camera matrix equation $\mathbf{P} = \mathbf{K}\mathbf{R}[\mathbf{I} \mid \mathbf{C}]$. Below the equation, four arrows point from labels to the terms \mathbf{K} , \mathbf{R} , $[\mathbf{I} \mid \mathbf{C}]$, and \mathbf{C} respectively. The labels are: 3×3 intrinsics, 3×3 3D rotation, 3×3 identity, and 3×1 3D translation.

3×3 intrinsics 3×3 3D rotation 3×3 identity 3×1 3D translation

Quiz

The camera matrix relates what two quantities?

Quiz

The camera matrix relates what two quantities?

$$\mathbf{x} = \mathbf{P}\mathbf{X}$$

3D world points to 2D image points, in homogeneous coordinates

Quiz

The camera matrix relates what two quantities?

$$\mathbf{x} = \mathbf{P}\mathbf{X}$$

3D world points to 2D image points, in homogeneous coordinates

The camera matrix can be decomposed into?

Quiz

The camera matrix relates what two quantities?

$$\mathbf{x} = \mathbf{P}\mathbf{X}$$

3D world points to 2D image points, in homogeneous coordinates

The camera matrix can be decomposed into?

$$\mathbf{P} = \mathbf{K}[\mathbf{R}|\mathbf{t}]$$


intrinsic and extrinsic parameters



Geometric camera calibration

(a.k.a. camera pose estimation)

Pose Estimation



Given a single image,
estimate the exact position of the photographer

Geometric camera calibration

Given a set of matched points

$$\{\mathbf{X}_i, \mathbf{x}_i\}$$

point in 3D
space

point in the
image

and camera model

$$\mathbf{x} = \mathbf{f}(\mathbf{X}; \mathbf{p}) = \mathbf{P}\mathbf{X}$$

projection
model

parameters

Camera
matrix

Find the (pose) estimate of

P

Mapping between 3D point and image points

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} p_1 & p_2 & p_3 & p_4 \\ p_5 & p_6 & p_7 & p_8 \\ p_9 & p_{10} & p_{11} & p_{12} \end{bmatrix} \begin{bmatrix} X \\ Y \\ Z \\ 1 \end{bmatrix}$$

What are the unknowns?

Mapping between 3D point and image points

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} p_1 & p_2 & p_3 & p_4 \\ p_5 & p_6 & p_7 & p_8 \\ p_9 & p_{10} & p_{11} & p_{12} \end{bmatrix} \begin{bmatrix} X \\ Y \\ Z \\ 1 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} \text{---} & \mathbf{p}_1^\top & \text{---} \\ \text{---} & \mathbf{p}_2^\top & \text{---} \\ \text{---} & \mathbf{p}_3^\top & \text{---} \end{bmatrix} \begin{bmatrix} | \\ \mathbf{X} \\ | \end{bmatrix}$$

Heterogeneous coordinates

$$x' = \frac{\mathbf{p}_1^\top \mathbf{X}}{\mathbf{p}_3^\top \mathbf{X}} \quad y' = \frac{\mathbf{p}_2^\top \mathbf{X}}{\mathbf{p}_3^\top \mathbf{X}}$$

(non-linear relation between coordinates)

How can we make these relations linear?

How can we make these relations linear?

$$x' = \frac{\mathbf{p}_1^\top \mathbf{X}}{\mathbf{p}_3^\top \mathbf{X}} \quad y' = \frac{\mathbf{p}_2^\top \mathbf{X}}{\mathbf{p}_3^\top \mathbf{X}}$$

Make them linear with algebraic manipulation...

$$\mathbf{p}_2^\top \mathbf{X} - \mathbf{p}_3^\top \mathbf{X} y' = 0$$

$$\mathbf{p}_1^\top \mathbf{X} - \mathbf{p}_3^\top \mathbf{X} x' = 0$$

Now we can setup a system of linear equations
with multiple point correspondences

$$\mathbf{p}_2^\top \mathbf{X} - \mathbf{p}_3^\top \mathbf{X} y' = 0$$

$$\mathbf{p}_1^\top \mathbf{X} - \mathbf{p}_3^\top \mathbf{X} x' = 0$$

How do we proceed?

$$\mathbf{p}_2^\top \mathbf{X} - \mathbf{p}_3^\top \mathbf{X} y' = 0$$

$$\mathbf{p}_1^\top \mathbf{X} - \mathbf{p}_3^\top \mathbf{X} x' = 0$$

In matrix form ...

$$\begin{bmatrix} \mathbf{X}^\top & \mathbf{0} & -x' \mathbf{X}^\top \\ \mathbf{0} & \mathbf{X}^\top & -y' \mathbf{X}^\top \end{bmatrix} \begin{bmatrix} \mathbf{p}_1 \\ \mathbf{p}_2 \\ \mathbf{p}_3 \end{bmatrix} = \mathbf{0}$$

How do we proceed?

$$\mathbf{p}_2^\top \mathbf{X} - \mathbf{p}_3^\top \mathbf{X} y' = 0$$

$$\mathbf{p}_1^\top \mathbf{X} - \mathbf{p}_3^\top \mathbf{X} x' = 0$$

In matrix form ...

$$\begin{bmatrix} \mathbf{X}^\top & \mathbf{0} & -x' \mathbf{X}^\top \\ \mathbf{0} & \mathbf{X}^\top & -y' \mathbf{X}^\top \end{bmatrix} \begin{bmatrix} \mathbf{p}_1 \\ \mathbf{p}_2 \\ \mathbf{p}_3 \end{bmatrix} = \mathbf{0}$$

For N points ...

$$\begin{bmatrix} \mathbf{X}_1^\top & \mathbf{0} & -x' \mathbf{X}_1^\top \\ \mathbf{0} & \mathbf{X}_1^\top & -y' \mathbf{X}_1^\top \\ \vdots & \vdots & \vdots \\ \mathbf{X}_N^\top & \mathbf{0} & -x' \mathbf{X}_N^\top \\ \mathbf{0} & \mathbf{X}_N^\top & -y' \mathbf{X}_N^\top \end{bmatrix} \begin{bmatrix} \mathbf{p}_1 \\ \mathbf{p}_2 \\ \mathbf{p}_3 \end{bmatrix} = \mathbf{0}$$

*How do we solve
this system?*

Solve for camera matrix by

$$\hat{\mathbf{x}} = \arg \min_{\mathbf{x}} \|\mathbf{A}\mathbf{x}\|^2 \text{ subject to } \|\mathbf{x}\|^2 = 1$$

$$\mathbf{A} = \begin{bmatrix} \mathbf{X}_1^\top & \mathbf{0} & -x' \mathbf{X}_1^\top \\ \mathbf{0} & \mathbf{X}_1^\top & -y' \mathbf{X}_1^\top \\ \vdots & \vdots & \vdots \\ \mathbf{X}_N^\top & \mathbf{0} & -x' \mathbf{X}_N^\top \\ \mathbf{0} & \mathbf{X}_N^\top & -y' \mathbf{X}_N^\top \end{bmatrix} \quad \mathbf{x} = \begin{bmatrix} p_1 \\ p_2 \\ p_3 \end{bmatrix}$$

SVD!

Solve for camera matrix by

$$\hat{\mathbf{x}} = \arg \min_{\mathbf{x}} \|\mathbf{A}\mathbf{x}\|^2 \text{ subject to } \|\mathbf{x}\|^2 = 1$$

$$\mathbf{A} = \begin{bmatrix} \mathbf{X}_1^\top & \mathbf{0} & -x' \mathbf{X}_1^\top \\ \mathbf{0} & \mathbf{X}_1^\top & -y' \mathbf{X}_1^\top \\ \vdots & \vdots & \vdots \\ \mathbf{X}_N^\top & \mathbf{0} & -x' \mathbf{X}_N^\top \\ \mathbf{0} & \mathbf{X}_N^\top & -y' \mathbf{X}_N^\top \end{bmatrix} \quad \mathbf{x} = \begin{bmatrix} p_1 \\ p_2 \\ p_3 \end{bmatrix}$$

Solution \mathbf{x} is the column of \mathbf{V}
corresponding to smallest singular
value of

$$\mathbf{A} = \mathbf{U}\mathbf{\Sigma}\mathbf{V}^\top$$

Solve for camera matrix by

$$\hat{\mathbf{x}} = \arg \min_{\mathbf{x}} \|\mathbf{A}\mathbf{x}\|^2 \text{ subject to } \|\mathbf{x}\|^2 = 1$$

$$\mathbf{A} = \begin{bmatrix} \mathbf{X}_1^\top & \mathbf{0} & -x' \mathbf{X}_1^\top \\ \mathbf{0} & \mathbf{X}_1^\top & -y' \mathbf{X}_1^\top \\ \vdots & \vdots & \vdots \\ \mathbf{X}_N^\top & \mathbf{0} & -x' \mathbf{X}_N^\top \\ \mathbf{0} & \mathbf{X}_N^\top & -y' \mathbf{X}_N^\top \end{bmatrix} \quad \mathbf{x} = \begin{bmatrix} p_1 \\ p_2 \\ p_3 \end{bmatrix}$$

Equivalently, solution \mathbf{x} is the Eigenvector corresponding to smallest Eigenvalue of

$$\mathbf{A}^\top \mathbf{A}$$

Now we have:

$$\mathbf{P} = \begin{bmatrix} p_1 & p_2 & p_3 & p_4 \\ p_5 & p_6 & p_7 & p_8 \\ p_9 & p_{10} & p_{11} & p_{12} \end{bmatrix}$$

Are we done?

Almost there ...

$$\mathbf{P} = \begin{bmatrix} p_1 & p_2 & p_3 & p_4 \\ p_5 & p_6 & p_7 & p_8 \\ p_9 & p_{10} & p_{11} & p_{12} \end{bmatrix}$$

How do you get the intrinsic and extrinsic parameters from the projection matrix?

Decomposition of the Camera Matrix

$$\mathbf{P} = \left[\begin{array}{ccc|c} p_1 & p_2 & p_3 & p_4 \\ p_5 & p_6 & p_7 & p_8 \\ p_9 & p_{10} & p_{11} & p_{12} \end{array} \right]$$

Decomposition of the Camera Matrix

$$\mathbf{P} = \left[\begin{array}{ccc|c} p_1 & p_2 & p_3 & p_4 \\ p_5 & p_6 & p_7 & p_8 \\ p_9 & p_{10} & p_{11} & p_{12} \end{array} \right]$$

$$\mathbf{P} = \mathbf{K}[\mathbf{R}|\mathbf{t}]$$

Decomposition of the Camera Matrix

$$\mathbf{P} = \left[\begin{array}{ccc|c} p_1 & p_2 & p_3 & p_4 \\ p_5 & p_6 & p_7 & p_8 \\ p_9 & p_{10} & p_{11} & p_{12} \end{array} \right]$$

$$\begin{aligned} \mathbf{P} &= \mathbf{K}[\mathbf{R}|\mathbf{t}] \\ &= \mathbf{K}[\mathbf{R} | -\mathbf{R}\mathbf{c}] \\ &= [\mathbf{M} | -\mathbf{M}\mathbf{c}] \end{aligned}$$

Decomposition of the Camera Matrix

$$\mathbf{P} = \left[\begin{array}{ccc|c} p_1 & p_2 & p_3 & p_4 \\ p_5 & p_6 & p_7 & p_8 \\ p_9 & p_{10} & p_{11} & p_{12} \end{array} \right]$$

$$\begin{aligned} \mathbf{P} &= \mathbf{K}[\mathbf{R}|\mathbf{t}] \\ &= \mathbf{K}[\mathbf{R} | -\mathbf{R}\mathbf{c}] \\ &= [\mathbf{M} | -\mathbf{M}\mathbf{c}] \end{aligned}$$

Find the camera center \mathbf{C}

What is the projection of the camera center?

Find intrinsic \mathbf{K} and rotation \mathbf{R}

Decomposition of the Camera Matrix

$$\mathbf{P} = \left[\begin{array}{ccc|c} p_1 & p_2 & p_3 & p_4 \\ p_5 & p_6 & p_7 & p_8 \\ p_9 & p_{10} & p_{11} & p_{12} \end{array} \right]$$

$$\begin{aligned} \mathbf{P} &= \mathbf{K}[\mathbf{R}|\mathbf{t}] \\ &= \mathbf{K}[\mathbf{R} | -\mathbf{R}\mathbf{c}] \\ &= [\mathbf{M} | -\mathbf{M}\mathbf{c}] \end{aligned}$$

Find the camera center \mathbf{c}

$$\mathbf{P}\mathbf{c} = \mathbf{0}$$

How do we compute the camera center from this?

Find intrinsic \mathbf{K} and rotation \mathbf{R}

Decomposition of the Camera Matrix

$$\mathbf{P} = \left[\begin{array}{ccc|c} p_1 & p_2 & p_3 & p_4 \\ p_5 & p_6 & p_7 & p_8 \\ p_9 & p_{10} & p_{11} & p_{12} \end{array} \right]$$

$$\begin{aligned} \mathbf{P} &= \mathbf{K}[\mathbf{R}|\mathbf{t}] \\ &= \mathbf{K}[\mathbf{R} | -\mathbf{R}\mathbf{c}] \\ &= [\mathbf{M} | -\mathbf{M}\mathbf{c}] \end{aligned}$$

Find the camera center \mathbf{c}

$$\mathbf{P}\mathbf{c} = \mathbf{0}$$

SVD of \mathbf{P} !

*\mathbf{c} is the Eigenvector corresponding to
smallest Eigenvalue*

Find intrinsic \mathbf{K} and rotation \mathbf{R}

Decomposition of the Camera Matrix

$$\mathbf{P} = \left[\begin{array}{ccc|c} p_1 & p_2 & p_3 & p_4 \\ p_5 & p_6 & p_7 & p_8 \\ p_9 & p_{10} & p_{11} & p_{12} \end{array} \right]$$

$$\begin{aligned} \mathbf{P} &= \mathbf{K}[\mathbf{R}|\mathbf{t}] \\ &= \mathbf{K}[\mathbf{R} | -\mathbf{R}\mathbf{c}] \\ &= [\mathbf{M} | -\mathbf{M}\mathbf{c}] \end{aligned}$$

Find the camera center \mathbf{c}

$$\mathbf{P}\mathbf{c} = \mathbf{0}$$

SVD of P!

\mathbf{c} is the Eigenvector corresponding to
smallest Eigenvalue

Find intrinsic \mathbf{K} and rotation \mathbf{R}

$$\mathbf{M} = \mathbf{K}\mathbf{R}$$

*Any useful properties of K
and R we can use?*

Decomposition of the Camera Matrix

$$\mathbf{P} = \left[\begin{array}{ccc|c} p_1 & p_2 & p_3 & p_4 \\ p_5 & p_6 & p_7 & p_8 \\ p_9 & p_{10} & p_{11} & p_{12} \end{array} \right]$$

$$\begin{aligned} \mathbf{P} &= \mathbf{K}[\mathbf{R}|\mathbf{t}] \\ &= \mathbf{K}[\mathbf{R} | -\mathbf{R}\mathbf{c}] \\ &= [\mathbf{M} | -\mathbf{M}\mathbf{c}] \end{aligned}$$

Find the camera center \mathbf{c}

$$\mathbf{P}\mathbf{c} = \mathbf{0}$$

SVD of P!

\mathbf{c} is the Eigenvector corresponding to
smallest Eigenvalue

Find intrinsic \mathbf{K} and rotation \mathbf{R}

$$\mathbf{M} = \mathbf{K}\mathbf{R}$$

↑ ↑
right upper orthogonal
triangle

*How do we find K
and R?*

Decomposition of the Camera Matrix

$$\mathbf{P} = \left[\begin{array}{ccc|c} p_1 & p_2 & p_3 & p_4 \\ p_5 & p_6 & p_7 & p_8 \\ p_9 & p_{10} & p_{11} & p_{12} \end{array} \right]$$

$$\begin{aligned} \mathbf{P} &= \mathbf{K}[\mathbf{R}|\mathbf{t}] \\ &= \mathbf{K}[\mathbf{R} | -\mathbf{R}\mathbf{c}] \\ &= [\mathbf{M} | -\mathbf{M}\mathbf{c}] \end{aligned}$$

Find the camera center \mathbf{c}

$$\mathbf{P}\mathbf{c} = \mathbf{0}$$

SVD of P!

\mathbf{c} is the Eigenvector corresponding to smallest Eigenvalue

Find intrinsic \mathbf{K} and rotation \mathbf{R}

$$\mathbf{M} = \mathbf{K}\mathbf{R}$$

QR decomposition

Geometric camera calibration

Given a set of matched points

$$\{\mathbf{X}_i, \mathbf{x}_i\}$$

point in 3D
space point in the
image

*Where do we get these
matched points from?*

and camera model

$$\mathbf{x} = \mathbf{f}(\mathbf{X}; \mathbf{p}) = \mathbf{P}\mathbf{X}$$

projection
model

parameters

Camera
matrix

Find the (pose) estimate of

P


Calibration using a reference object

Place a known object in the scene:

- identify correspondences between image and scene
- compute mapping from scene to image

Issues:

- must know geometry very accurately
- must know 3D->2D correspondence



Geometric camera calibration

Advantages:

- Very simple to formulate.
- Analytical solution.


Disadvantages:

- Doesn't model radial distortion.
- Hard to impose constraints (e.g., known f).
- Doesn't minimize the correct error function.

For these reasons, *nonlinear methods* are preferred

- Define error function E between projected 3D points and image positions
 - E is nonlinear function of intrinsics, extrinsics, radial distortion
- Minimize E using nonlinear optimization techniques

Known 2d image coords



880 214
43 203
270 197
886 347
745 302
943 128
476 590
419 214
317 335
783 521
235 427
665 429
655 362
427 333
412 415
746 351
434 415
525 234
716 308
602 187

Known 3d world locations

312.747 309.140 30.086
305.796 311.649 30.356
307.694 312.358 30.418
310.149 307.186 29.298
311.937 310.105 29.216
311.202 307.572 30.682
307.106 306.876 28.660
309.317 312.490 30.230
307.435 310.151 29.318
308.253 306.300 28.881
306.650 309.301 28.905
308.069 306.831 29.189
309.671 308.834 29.029
308.255 309.955 29.267
307.546 308.613 28.963
311.036 309.206 28.913
307.518 308.175 29.069
309.950 311.262 29.990
312.160 310.772 29.080
311.988 312.709 30.514

Known 2d
image coords
(px)


Known 3d
world locations
(m)

$$\begin{bmatrix} su \\ sv \\ s \end{bmatrix} = \begin{bmatrix} m_{11} & m_{12} & m_{13} & m_{14} \\ m_{21} & m_{22} & m_{23} & m_{24} \\ m_{31} & m_{32} & m_{33} & m_{34} \end{bmatrix} \begin{bmatrix} X \\ Y \\ Z \\ 1 \end{bmatrix}$$


M

Unknown Camera Parameters

3D locations of planar marker features are known in advance



3D content prepared in advance



Simple AR program

1. Compute point correspondences (2D and AR tag)
2. Estimate the pose of the camera **P**
3. Project 3D content to image plane using **P**

More Advance Calibration using Multiple Views....





<https://grandvisual.com/work/pepsi-max-bus-shelter/> (London, 2014)



The University of Texas at Austin
**Electrical and Computer
Engineering**
Cockrell School of Engineering